

Conics in the Grothendieck ring

János Kollár

February 8, 2008

The aim of this note is to describe the subring of the Grothendieck ring generated by smooth conics. As a ring this is quite complicated, with many zero divisors, but the description of the defining relations is entirely elementary.

Definition 1. Let k be a field. The *Grothendieck ring* of k -varieties, denoted by $K_0[\mathrm{Var}_k]$ is defined as follows.

Its additive group is the Abelian group whose generators are the isomorphism classes of reduced, quasi projective k -schemes and the relations are

$$[X] = [Y] + [X \setminus Y],$$

whenever Y is a closed subscheme of X .

Multiplication is defined by $[X] \cdot [Y] = [X \times_k Y]$.

The Grothendieck ring of k -varieties is still very poorly understood. In characteristic zero, the quotient of $K_0[\mathrm{Var}_k]$ by the ideal generated by $[\mathbb{A}^1]$ is naturally isomorphic to the ring $\mathbb{Z}[SB_k]$, where $\mathbb{Z}[SB_k]$ is the free abelian group generated by the stable birational equivalence classes of smooth, projective, irreducible k -varieties and multiplication is given by the product of varieties [Lar-Lun]. (The cited paper proves this over algebraically closed fields only, but the proof works over any field of characteristic zero using the birational factorization theorem as given in [AKMW, Remark 2 after Theorem 0.3.1]. Note also that the product of two irreducible k -varieties is not necessarily irreducible, so $\mathbb{Z}[SB_k]$ is not a monoid ring if k is not algebraically closed.)

Zero divisors in the Grothendieck ring of $\mathbb{C}$ -varieties were found by [Poonen].

Here we give further examples of nontrivial behaviour of these rings by studying products of conics. This gives interesting examples only when the field k is not algebraically closed.

Theorem 2. *Let k be a number field or the function field of an algebraic surface over $\mathbb{C}$. Let $C_i : i \in I$ and $C'_j : j \in J$ be two collections of smooth conics defined over k (repetitions allowed).*

Then $[\prod_{i \in I} C_i] = [\prod_{j \in J} C'_j]$ in the Grothendieck ring iff $|I| = |J|$ and the subgroup of the Brauer group (cf. (7)) generated by the first collection $\langle C_i : i \in I \rangle \subset \mathrm{Br}(k)_2$ is the same as the subgroup of the Brauer group generated by the second collection $\langle C'_j : j \in J \rangle \subset \mathrm{Br}(k)_2$.

Remark 3. (1) The precise conditions on k for the proof to work are given in (8). These are satisfied for many other fields, but fail for function fields of more than 2 variables. It is not clear to me, however, if any condition is needed on k .

(2) Any isomorphism of two products $\prod_{i \in I} C_i$ and $\prod_{j \in J} C'_j$ is given in the obvious way: by a one-to-one map $g : I \rightarrow J$ and isomorphisms $C_i \cong C'_{g(i)}$ for every i .

This can be proved many ways. Here is one using extremal rays.

If X is any projective variety, the cone of curves of $X \times \mathbb{P}^1$ is generated by the cone of curves of $X \cong X \times \{0\}$ and by $\{x\} \times \mathbb{P}^1$. Using this repeatedly, we obtain that the cone of curves of $(\mathbb{P}^1)^m$ is generated by the fibers of the m coordinate projections $(\mathbb{P}^1)^m \rightarrow (\mathbb{P}^1)^{m-1}$. Thus the $|I|$ coordinate projections

$$\pi_{i'} : \prod_{i \in I} C_i \rightarrow \prod_{i \in I, i \neq i'} C_i$$

are in one-to-one correspondence with the extremal rays of $\prod_{i \in I} C_i$. Hence the product structure can be recovered from the intrinsic geometry of $\prod_{i \in I} C_i$.

Corollary 4. *Let k be a number field or the function field of an algebraic surface over $\mathbb{C}$. The subring of the Grothendieck ring generated by smooth conics is isomorphic to the ring generated by the isomorphism classes of smooth conics modulo the ideal generated by the relations $[C_1] \cdot [C_2] - [C_1] \cdot [C_1 * C_2]$.*

This ring can also be described as follows.

Let $G \subset \text{Br}(k)_2$ be a finite subgroup with basis $B_1, \dots, B_s$. Then $[B_1 \times \dots \times B_s]$ depends only on G and it is denoted by $C(G)$. The trivial subgroup gives $C(0) = [\text{Spec } k]$.

The Grothendieck ring of conics is the free abelian group generated by the elements $C(G) \cdot [\mathbb{P}^1]^m$ with multiplication

$$C(G_1) \cdot C(G_2) = C(\langle G_1, G_2 \rangle) \cdot [\mathbb{P}^1]^{\dim G_1 + \dim G_2 - \dim \langle G_1, G_2 \rangle}.$$

where $\dim G$ denotes dimension as an $\mathbb{F}_2$ vector space.

Remark 5. The last description shows that the Grothendieck ring of conics does not have nilpotents. Indeed, given an element $g = \sum \gamma_{G,m} C(G) \cdot [\mathbb{P}^1]^m$, let G_0 be minimal such that $\gamma_{G_0,m} \neq 0$ for some m , chosen also minimal. Then the coefficient of $C(G_0) \cdot [\mathbb{P}^1]^{sm + (s-1)\dim G_0}$ in g^s is $\gamma_{G_0,m}^s \neq 0$.

The simplest example of nontrivial birational maps between products of conics is the following. The whole description of the Grothendieck ring of conics is only a more elaborate version of it.

Example 6. Let C be a smooth plain conic. Then $C \times \mathbb{P}^1$ is birational to $C \times C$, and they have the same class in the Grothendieck ring of k -varieties.

Thus $[C] \cdot ([\mathbb{P}^1] - [C]) = 0$ and $[C]$ is a zero divisor in the Grothendieck ring of k -varieties if C has no k -points.

Proof. $C \subset \mathbb{P}^2$ is a conic and we think of $\mathbb{P}^1$ as a line in the same $\mathbb{P}^2$.

Given $p, q \in C$, the line connecting them intersects $\mathbb{P}^1$ in a point $\phi(p, q)$. $(p, q) \mapsto (p, \phi(p, q))$ gives a rational map $C \times C \dashrightarrow C \times \mathbb{P}^1$. Conversely, given $p \in C$ and $r \in \mathbb{P}^1$ the line connecting them intersects C in a further point $\phi^{-1}(p, r)$.

Let $s, s' \in C(\bar{k})$ be the two intersection points of C and $\mathbb{P}^1$. ϕ is not defined at the pairs (s, s') and (s', s) . ϕ^{-1} is not defined at the pairs (s, s) and (s', s') . Easy computation shows that ϕ becomes an isomorphism after we blow up the indeterminacy loci. The blown-up surface is denoted by $B(C \times C)$. As k -schemes, $(s, s') \cup (s', s)$ and $(s, s) \cup (s', s')$ are both isomorphic to $\text{Spec}_k k(s)$.

Thus $[C \times C]$ and $[C \times \mathbb{P}^1]$ can both be written as

$$[B(C \times C)] - [\mathbb{P}^1] \cdot [\text{Spec}_k k(s)] + [\text{Spec}_k k(s)].$$

In order to see that $[C]$ is a zero divisor in the Grothendieck ring of k -varieties, we need to prove that $[\mathbb{P}^1] - [C]$ is not zero. By [Lar-Lun], it is sufficient to prove that $\mathbb{P}^1$ and C are not stably birational. This is however easy, since having k -points is a stably birational invariant.

7 (Products of conics and the Brauer group).

Below we give an elementary geometric description of the Brauer group of conics, denoted by $\text{Br}(k)_2$. (For the fields that we are considering, this is the 2-torsion subgroup of the Brauer group $\text{Br}(k)$. See [Serre, X.4–7] for a good introduction and basic properties.)

Let k be a field and C_1, C_2 two smooth conics defined over k . The Brauer product of the two conics is defined as follows. (I warn the reader in advance that this definition only works because on a conic the Hilbert scheme of points is isomorphic to the Hilbert scheme of degree 1 divisors. On a higher dimensional Severi–Brauer variety the Hilbert scheme of points is dual to the Hilbert scheme of degree 1 divisors and one has to distinguish these systematically.)

Start with $C_1 \times C_2$. As a first approximation, we construct a 3-dimensional variety, denoted by $P(C_1, C_2)$. We would like to say that $P(C_1, C_2)$ is the 3-dimensional “linear system” of divisors of bidegree $(1, 1)$ on $C_1 \times C_2$. The problem is that in general no such divisor is defined over k . Thus we look at the linear system $|-K|$ where $K = K_{C_1 \times C_2}$ is the canonical class. This corresponds to divisors of bidegree $(2, 2)$. Then $P(C_1, C_2) \subset |-K|$ is the subscheme consisting of those divisors which are everywhere double. Over the algebraic closure of k we recognize this as the (doubled) elements of the linear system $|\mathcal{O}(1, 1)|$.

Alternatively, the Hilbert scheme $\text{Hilb}(C_1 \times C_2)$ has an irreducible component parametrizing divisors of bidegree $(1, 1)$. This is again $P(C_1, C_2)$.

Thus $P(C_1, C_2)$ is a 3-dimensional k -variety which is isomorphic to $\mathbb{P}^3$ over $\bar{k}$.

There is a natural embedding $C_1 \times C_2 \hookrightarrow P(C_1, C_2)$ where we map a point $(p, q) \in C_1 \times C_2$ to the divisor $2(\{p\} \times C_2 + C_1 \times \{q\})$.

In general this is all one can do. There are, however, important cases when such a product $P(C_1, C_2)$ contains a degree 1 smooth curve (a line over $\bar{k}$) defined over k . In this case I call this degree 1 curve the Brauer product of C_1 and C_2 and denote it by $C_1 * C_2$. (The terminology “Brauer product” does not seem to be standard.)

It turns out that this is well defined.

To see this, let P be a 3-dimensional k -variety which is isomorphic to $\mathbb{P}^3$ over $\bar{k}$. Let $L_1, L_2 \subset P$ be degree 1 smooth curves defined over k and let $L' \subset P$ be another such curve disjoint from both. (Over an infinite field we can obtain L' as the image of L_1 by a general automorphism of P .) Then L_1 and L_2 are both isomorphic to the Hilbert scheme of degree 1 surfaces containing L' .

$C * C$ is always defined and it is isomorphic to $\mathbb{P}^1$. Indeed, the diagonal $\Delta \subset C \times C$ is defined over k thus $P(C, C)$ is k -isomorphic to $\mathbb{P}_k^3$. Hence the Brauer group of conics is a 2-group.

Lemma 8. *For a field k the following two conditions are equivalent.*

1. *The Brauer product of 2 smooth conics is again a conic.*
2. *For any two smooth conics C_1, C_2 defined over k there is a degree 2 extension k'/k such that both C_1 and C_2 have k' -points.*

Proof. Let $L \subset P(C_1, C_2)$ be a degree 1 curve defined over k . Then $L \cap (C_1 \times C_2)$ is a degree 2 subscheme defined over k with residue field k' . By projection to the factors, C_1, C_2 both have points in k' .

Conversely, if C_1, C_2 both have points in k' then so does their product. The unique line in $P(C_1, C_2)$ passing through a k' point is defined over k . $\square$

The following result is well known in various forms, see for instance [Artin, p.209] or [Sarkisov, Thm.5.7]

Proposition 9. *The conditions in (8) hold in the following two cases:*

1. *k is a number field.*
2. *k is the function field of an algebraic surface over an algebraically closed field. More generally, for C_2 -fields.*

Proof. Here is a geometric version of some of the classical proof.

Let $G(1, P(C_1, C_2))$ denote the Grassmannian of lines in $P(C_1, C_2)$. We need to prove that it has a k -point.

More generally, let P be a k -variety which is isomorphic to $\mathbb{P}^n$ over $\bar{k}$ and assume that there is a quadric hypersurface $Q \subset P$ defined over k . As explained in [Artin, 4.5] the Grassmannian of lines $G(1, P)$ is embedded into $\mathbb{P}^{\binom{n+1}{2}-1}$ the usual way.

For $n = 2$ the Grassmannian of lines $G(1, P)$ is thus a quadric in $\mathbb{P}^5$, and so it has a point over any C_2 field, proving the second part.

If $k = \mathbb{R}$ then a $\mathbb{C}$ -point of P and its conjugate determine a real line, so $G(1, P)(\mathbb{R}) \neq \emptyset$. Thus $G(1, P)$ is a quadric in 6 variables which has a point in all real completions of k . Therefore $G(1, P)$ has a k -point by the Hasse–Minkowski theorem. $\square$

10 (Proof of (2)).

The key point to show one direction is the following generalization of (6):

Lemma 11. *Let C_1, C_2 be smooth conics such that their Brauer product $C_1 * C_2$ is again a smooth conic. Then*

1. $C_1 \times C_2$ is birational to $C_1 \times (C_1 * C_2)$.
2. $[C_1 \times C_2] = [C_1 \times (C_1 * C_2)]$ in the Grothendieck ring.

Proof. First we write down a rational map $\phi : C_1 \times C_2 \dashrightarrow C_1 * C_2$. Then we check that ϕ and the first projection $\pi_1 : C_1 \times C_2 \rightarrow C_1$ give a birational map

$$(\pi_1, \phi) : C_1 \times C_2 \dashrightarrow C_1 \times (C_1 * C_2).$$

Finally we see that this gives an identity in the Grothendieck ring.

Geometric description. By assumption there is a degree 2 point $Q \in C_1 \times C_2 \subset P(C_1, C_2)$. Let L' be the unique degree 1 curve through Q and let $L \in P(C_1, C_2)$ be any degree 1 curve disjoint from L' . Projection from L' to L gives ϕ .

Algebraic description. Assume for simplicity that the characteristic is different from 2. If the common point is in the field $k(\sqrt{a})$, we can assume that the conics are given by equations

$$C_1 = (x_1^2 - ax_2^2 - bx_3^2 = 0) \quad \text{and} \quad C_2 = (y_1^2 - ay_2^2 - cy_3^2 = 0).$$

Then their Brauer product can be given as

$$C_1 * C_2 = (z_1^2 - az_2^2 - bcz_3^2 = 0)$$

and ϕ is given by

$$(z_1 : z_2 : z_3) = (x_1y_1 + ax_2y_2 : x_1y_2 + x_2y_1 : x_3y_3).$$

ϕ^{-1} is obtained as follows. Pick a point $p \in C_1$ and $r \in L \cong C_1 * C_2$. $\{p\} \times C_2$ embeds as a line into $P(C_1, C_2)$ and $\phi^{-1}(p, r)$ is the intersection point of this line with the plane $\langle L', r \rangle$ spanned by L' and r . This is not defined only if $\{p\} \times C_2 \subset \langle L', r \rangle$. This happens exactly when $\langle L', r \rangle$ is one of the two tangent planes of $C_1 \times C_2$ at a point of Q and $\{p\} \times C_2$ is the corresponding line through that point of Q .

Thus we see that $C_1 \times C_2$ becomes isomorphic to $C_1 \times (C_1 * C_2)$ after we blow up subschemes isomorphic to Q in both of them. As in (6) this shows that $[C_1 \times C_2] = [C_1 \times (C_1 * C_2)]$. $\square$

Assume that the subgroup $G_I \subset \text{Br}(k)_2$ generated by the C_i -s is the same as the subgroup $G_J \subset \text{Br}(k)_2$ generated by the C'_j -s. Fix a minimal generating set $\{B_s : s \in S\}$ of G . By a simple group theoretic lemma (12) and a repeated application of (11), $\prod_{i \in I} C_i$ is birational to

$$(\mathbb{P}_k^1)^{|I|-|S|} \times \prod_{s \in S} B_s,$$

and they have the same class in the Grothendieck ring. The same holds for $\prod_{j \in J} C'_j$. Thus $\prod_{i \in I} C_i$ and $\prod_{j \in J} C'_j$ are birational and they have the same class in the Grothendieck ring.

Conversely, assume that $G_I \neq G_J$. We may assume that $G_J \not\subset G_I$ and so there is an index j_0 such that the class of C'_{j_0} is not in G_I . We claim that in this case there is no rational map from $\prod_{i \in I} C_i$ to C'_{j_0} , hence no rational map from $\prod_{i \in I} C_i$ to $\prod_{j \in J} C'_j$. Thus they are not birational and not even stably birational, hence they represent different elements of the Grothendieck ring by [Lar-Lun].

The proof is by induction on $|I|$, the case $|I| = 0$ being clear. Pick $i_0 \in I$ and set $I' := I \setminus \{i_0\}$ and $K = k(C_{i_0})$. By (13), the kernel of $G_I \rightarrow \text{Br}(k)_2$ is generated by C_{i_0} and so the class of C'_{j_0} in $\text{Br}(k)_2$ is not in the subgroup $G_{I'} \subset \text{Br}(K)_2$ generated by the C_i -s for $i \in I'$. By induction, there is no $k(C_{i_0})$ -map from $\prod_{i \in I'} C_i$ to C'_{j_0} , and so no k -map from $\prod_{i \in I} C_i$ to C'_{j_0} . This completes the proof of (2).

In the proof we used only the relations given by (11), and this gives the first description of the Grothendieck ring in (4).

It is clear the all the possible products $\prod_{i \in I} C_i$ generate the Grothendieck ring as an additive abelian group, and we have established that each such product is identical to a unique element of the form

$$[\mathbb{P}_k^1]^{|I|-|S|} \cdot \prod_{s \in S} [B_s] = [\mathbb{P}_k^1]^{|I|-|S|} \cdot C(G).$$

This gives the second description in (4). $\square$

Lemma 12. *Let G be a finite abelian 2-group with a minimal generating set $b_1, \dots, b_m$.*

Let $e_1, \dots, e_s$ be any generating collection of elements of G , repetitions allowed. Then $e_1, \dots, e_s$ can be transformed into the collection $b_1, \dots, b_m, 0, \dots, 0$ by repeated application of the following operation:

Pick e_i, e_j and replace e_j by $e_i + e_j$. $\square$

The following is a very special case of an old result of [Amitsur].

Lemma 13. *Let C, C' be smooth conics and $g : C \rightarrow C'$ a rational map. Then either $C' \cong \mathbb{P}^1$ or $C \cong C'$.*

Therefore, if $C' \not\cong C, \mathbb{P}^1$ then C' does not have a $k(C)$ -point.

Proof. Let $G \subset C \times C'$ be the graph of g . It is a divisor of bidegree $(1, \deg g)$. A class of bidegree $(0, 2)$ is defined over k , so we obtain that either the linear system $|\mathcal{O}(1, 0)|$ or the linear system $|\mathcal{O}(1, 1)|$ has a member over k . In the first case C' has a k -point and $C' \cong \mathbb{P}^1$ and in the second case we get a graph of an isomorphism. $\square$

The birational part of (2) is easy to state and prove for higher dimensional Severi–Brauer varieties:

Proposition 14. *Two products $\prod_{i \in I} P_i$ and $\prod_{j \in J} P'_j$ of Severi–Brauer varieties are stably birational iff the subgroup $\langle P_i : i \in I \rangle \subset \text{Br}(k)$ is the same as the subgroup $\langle P'_j : j \in J \rangle \subset \text{Br}(k)$.*

It is less clear how to formulate the Grothendieck ring version in general. Even for products of 2–dimensional Severi–Brauer varieties I found it difficult to write down a suitable analog of (11). Additional problems arise when the two Severi–Brauer varieties do not have a common splitting field.

Acknowledgments . Partial financial support was provided by the NSF under grant number DMS02-00883.

References

- [AKMW] D. Abramovich, K. Karu, K. Matsuki and J. Włodarczyk, Torification and factorization of birational maps. J. Amer. Math. Soc. 15 (2002), no. 3, 531–572
- [Amitsur] S. Amitsur, Generic splitting fields of central simple algebras, Ann. Math. 62 (1955) 8–43
- [Artin] M. Artin, Brauer-Severi varieties. in: Brauer groups in ring theory and algebraic geometry (Wilrijk, 1981), 194–210, Lecture Notes in Math. 917, Springer, 1982.
- [Lar-Lun] M. Larsen and V. Lunts. Motivic measures and stable birational geometry. math.AG/0110255
- [Poonen] B. Poonen, The Grothendieck ring of varieties is not a domain. math.AG/0204306
- [Sarkisov] V. G. Sarkisov, On conic bundle structures. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 46 (1982), no. 2, 371–408
- [Serre] J.-P. Serre, Local fields, Graduate Texts in Mathematics, 67. Springer-Verlag, New York-Berlin, 1979

Princeton University, Princeton NJ 08544-1000

`kollar@math.princeton.edu`